



Women's Health Disparities

Adopted by the SOPHE Board of Trustees

May 12, 2021

Whereas, more than 50% of the United States (US) population are female (US Census Bureau [USCB], n.d.a) with 76% identifying as white, 13% as African American, 6% as Asian, and 18% Hispanic (USCB, n.d.b).

Whereas, more than 13% of women 18 years of age and older in 2018 were in poor or fair health (US Department of Health and Human Services [HHS], 2018).

Whereas, women are twice as likely to be diagnosed with depression, Post Traumatic Stress Disorder, (PTSD), and experience anxiety (Kuehner, 2016; Office of Women's Health, 2019; U.S. Department of Veterans Affairs, 2018).

Whereas, more than 41 million US women are annually diagnosed with cancer (Centers for Disease Control and Prevention [CDC], 2016).

Whereas, almost 30% of US women over 50 report never having a colorectal cancer screening (Kaiser Family Foundation [KFF], 2019a).

Whereas, 1 in 3 US women report not having a dental exam in the past year (KFF, 2019b).

Whereas, women and girls report engaging in less physical activity (Armstrong et al., 2018; Telford et al., 2016).

Whereas, 15 million women are diagnosed with diabetes annually (CDC, 2018). Diabetes increases the risk of heart disease in women by four times. Women experience worse outcomes following a heart attack, and women are at a higher risk for experiencing diabetes-related complications (Centers for Disease Control and Prevention, 2017; Centers for Disease Control & Prevention, 2018).

Whereas, women have a higher lifetime risk of stroke when compared to men, and they comprise a larger portion of deaths due to stroke. Among all women, African American women are more likely to have a stroke (Benjamin et al., 2019).

Whereas, health disparities exist between women based on race, ethnicity, geography, and age (KFF, 2019c).

Whereas, Non-Hispanic Black women (ages 15-24 years), are more likely to be impacted by homicide (37.2%) and heart disease (11%) compared to non-Hispanic White women (8.7% and 6%, respectively) (Heron, 2019).

Whereas, among all US women, 17% report not having a primary care physician; Hispanic women, Asian women, and Native American women reporting higher instances at 32%, 24%, and 19% respectively (KFF, 2019d).

Whereas, 15% of adult US women don't see a doctor because of cost with this rate being higher among African American (19%), Hispanic (23%), and Native American (21%) women (KFF, 2019e).

Whereas, among US women over 40 years, 27% have not had a mammogram in the past two years with 33% of Native American reporting an even higher percentage (33%) (KFF, 2019f).

Whereas, health disparities are worse in the Southern states as non-elderly adult women face higher uninsured rates TX (22%), OK (19%), MS (17%), GA (17%), AL (13%), FL (17%), SC (15%), NC (14%) (KFF, 2019g).

Whereas, in 2017, women between the ages of 19 and 64 years accessed health insurance in a variety of ways: 60% of women received health care coverage through employers, 13% received coverage via non-group or other public insurance, and 17% received coverage through Medicaid, yet, 11% were uninsured (Kaiser Family Foundation [KFF], 2018).

Whereas, in 2014, women comprised 36% of the population receiving Medicaid, yet men comprised 21% of the population receiving Medicaid (Kaiser Family Foundation [KFF], 2019h).

Whereas, among LGBT individuals, disparities exist for Lesbian and Bisexual women as they are more likely to be overweight or obese (Struble et al., 2010; U.S. Department of Health and Human Services, 2019). Women who are Lesbians are also less likely to obtain preventive services for cancer (Buchmueller & Carpenter, 2010; Dilley et al., 2010).

Whereas, among clinical trials for heart failure, coronary artery disease, and acute coronary syndrome, women remain underrepresented (Scott et al., 2018).

Therefore, be it resolved, the Society for Public Health Education, Inc. (SOPHE) shall:

Urge Congress to:

- A. Advocate for:
 - a. funding to decrease health disparities between and among women
 - b. affordable healthcare that includes coverage for preventative care
 - c. funding to increase participation in clinical trials among women
 - d. allocate additional funding to the National Institutes of Health Office of Research on Women's Health

- B. Support and advocate for implementation of evidence-based programs and activities that increase health equity and reduce health disparities in women’s health.
- C. Support policies that are driven by science-based research.
- D. Oppose political interference between physicians/health providers and their patients.
- E. Support the Equality Act or other legislation coming forward that support LGBTQ women have access to explicit and comprehensive nondiscrimination protections.

Encourage and support efforts within public health to:

- A. Support local communities in the delivery of social services focused on women’s health.
- B. Support local communities to promote safe schools, neighborhoods, and housing for all women.
- C. Continue collecting data to identify trends and health disparities to address specific women’s health concerns.
- D. Collaborate with other organizations (National, State, and Community) to work together to:
 - a. Reduce women’s health disparities
 - b. Increase access to preventative services for all women

Internal Activities (for SOPHE, SOPHE Chapters, and Members)

1. Provide trainings, conference sessions, and pre-conference workshops focusing on issues related to women’s health disparities.
2. Update materials, trainings, and workshops to include inclusive language and be culturally inclusive of all women.
3. Demonstrate public support for women’s health and health care access for all women.
4. Promote inclusivity by doing activities such as ensuring speakers are culturally inclusive and using inclusive language, seeking out opportunities for venues that support diversity, continuing to provide examples of individuals or organizations who promote inclusivity of all women.

References

- Armstrong, S., Wong, C. A., Perrin, E., Page, S., Sibley, L., & Skinner, A. (2018). Association of physical activity with income, race/ethnicity, and sex among adolescents and young adults in the United States. *JAMA Pediatrics*, 172(8), 732-740. doi:10.1001/jamapediatrics.2018.1273
- Benjamin, E. J., Muntner, P., Alonso, A., Bittencourt, M. S., Callaway, C. W., Carson, A. P...Virani, S. S. (2019). Heart disease and stroke statistics – 2019 update: A report from the American Heart Association. *Circulation*, 139, e56-e528
- Buchmueller, T. & Carpenter, C.S. (2016). Disparities in health insurance coverage, access, and outcomes for individuals in same-sex versus different-sex relationships, 2000–2007. *American Journal of Public Health*, 100(3):489-95.
- Centers for Disease Control and Prevention. (2016). United States Cancer Statistics. Retrieved from <https://gis.cdc.gov/Cancer/USCS/DataViz.html>

- Centers for Disease Control and Prevention. (2017). *National diabetes statistics report, 2017*. Atlanta, GA: Centers for Disease Control and Prevention, U.S. Department of Health and Human Services.
- Centers for Disease Control and Prevention. (2018). Diabetes and women. Retrieved from <https://www.cdc.gov/features/diabetes-women/index.html>.
- Dilley, J.A, Simmons, K.W., Boysun, M.J., et al. (2010). Demonstrating the importance and feasibility of including sexual orientation in public health surveys: Health disparities in the Pacific Northwest. *American Journal of Public Health*, 100(3), 460-7.
- Kaiser Family Foundation. (2019a). *Percent of Women age 50 or Older who Report Ever Having Had a Colorectal Cancer Screening*. Retrieved from <https://www.kff.org/womens-health-policy/state-indicator/colorectal-cancer-screening-rate/?currentTimeframe=0&sortModel=%7B%22colld%22:%22Location%22,%22sort%22:%22asc%22%7D>
- Kaiser Family Foundation. (2019b). *Percent of Women who Visited a Dentist or Dental Clinic within the Past Year*. Retrieved from <https://www.kff.org/womens-health-policy/state-indicator/percent-of-women-who-had-dental-visit/?currentTimeframe=0&sortModel=%7B%22colld%22:%22Location%22,%22sort%22:%22asc%22%7D>
- Kaiser Family Foundation. (2019c). *Minority Health Indicators*. Retrieved from <https://www.kff.org/state-category/minority-health/womens-health-disparities/>
- Kaiser Family Foundation. (2019d). *Percent of Women who Report Having no Personal Doctor/Health Care Provider, by Race/Ethnicity*. Retrieved from <https://www.kff.org/disparities-policy/state-indicator/no-personal-doctor/?currentTimeframe=0&sortModel=%7B%22colld%22:%22Location%22,%22sort%22:%22asc%22%7D>
- Kaiser Family Foundation. (2019e). *Percent of Adult Women who did not see a Doctor in the Past 12 Months Due to Cost, by Race/Ethnicity*. Retrieved from <https://www.kff.org/womens-health-policy/state-indicator/percent-of-adult-women-who-did-not-see-a-doctor-in-the-past-12-months-due-to-cost-by-raceethnicity/?currentTimeframe=0&sortModel=%7B%22colld%22:%22Location%22,%22sort%22:%22asc%22%7D>
- Kaiser Family Foundation. (2019f). *Percent of Women age 40 and Older who Report Having Had a Mammogram Within the Past Two Years, by Race/Ethnicity*. Retrieved from <https://www.kff.org/womens-health-policy/state-indicator/mammogram-rate-for-women-40-years/?currentTimeframe=0&sortModel=%7B%22colld%22:%22Location%22,%22sort%22:%22asc%22%7D>
- Kaiser Family Foundation. (2019g). *Key Facts about the Uninsured Population*. Retrieved from <https://www.kff.org/uninsured/fact-sheet/key-facts-about-the-uninsured-population/>
- Kaiser Family Foundation [KFF]. (2019h). Health insurance coverage of women 19-64. Retrieved from <https://www.kff.org/other/state-indicator/nonelderly-adult-women/?currentTimeframe=0&sortModel=%7B%22colld%22:%22Other%20Public%22,%22sort%22:%22asc%22%7D>
- Kuehner, C. (2017). Why is depression more common among women than among men? *The Lancet Psychiatry*, 4(2), 146-158. doi: [https://doi.org/10.1016/S2215-0366\(16\)30263-2](https://doi.org/10.1016/S2215-0366(16)30263-2)
- Office of Women's Health. (2019). Anxiety disorders. Retrieved from <https://www.womenshealth.gov/mental-health/mental-health-conditions/anxiety-disorders#19>

- Scott, P. E., Unger, E. F., Jenkins, M. R., Southworth, M. R., McDowell, T., Geller, R. G., Elahi, M., Temple, R. J., & Woodcock, J. (2018). Participation of women in clinical trials supporting FDA approval of cardiovascular drugs. *Journal of the American College of Cardiology*, *71*(18), 1960-1969. doi:10.1016/j.jacc.2018.02.070
- Struble, C.B., Lindley, L.L., Montgomery, K., & Burcin, M. (2010). Overweight and obesity in lesbian and bisexual college women. *Journal of American College Health*, *59*(1), 51-6.
- Telford, R. M., Telford, R. D., Olive, L. S., Cochrane, T., & Davey, R. (2016). Why girls are less physically active than boys? Findings from the LOOK longitudinal study. *PLoS ONE*, *11*(3). doi:10.1371/journal.pone.0150041
- United States Department of Health and Human Services. (2018). Summary Health Statistics: National Health Interview Survey. Retrieved from https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-11.pdf
- U.S. Department of Health and Human Services. (2019). Healthy People 2020: Lesbian, gay, bisexual, and transgender health. Retrieved from <https://www.healthypeople.gov/2020/topics-objectives/topic/lesbian-gay-bisexual-and-transgender-health#22>
- U.S. Department of Veterans Affairs. (2018). PTSD: National center for PTSD. Retrieved from https://www.ptsd.va.gov/understand/common/common_women.asp.
- United States Census Bureau. (n.d.a). *Quick Facts, United States*. Retrieved from <https://www.census.gov/quickfacts/fact/table/US/LFE046217>
- United States Census Bureau. (n.d.b). *Quick Facts, United States, Female Persons*. Retrieved from <https://www.census.gov/quickfacts/fact/table/US/SEX255218>

Recommended Citation:

Society for Public Health Education. (2021). *Women's health disparities*. URL